



---

## Department of Physics, IIT-Kanpur

Time : 2 hrs.

PhD Admission Test May 2019

Total Marks: 70

### Question 1

Consider a free particle wave function in one dimension is given by

$$\psi(x) = A \exp\left\{-\frac{1}{2} \frac{(x-a)^2}{x_0^2}\right\},$$

where  $a$  and  $x_0$  are constants. Find  $\langle x \rangle$ ,  $\langle x^2 \rangle$ ,  $\langle p \rangle$ ,  $\langle p^2 \rangle$  and the uncertainty product. [5]

### Question 2

(a) Find the eigenvalues of the operator  $\hat{A} = 2\hat{J}_x + 2\hat{J}_y + \hat{J}_z$ , where the operators denote the components of the angular momentum operator with a magnitude  $J = 2$ . [2]

(b) Consider a state  $|\psi\rangle = a|2, 2\rangle + b|2, 0\rangle + c|2, -2\rangle$  written in  $|j, m\rangle$  basis. Find the expectation value  $\langle \psi | \hat{A} | \psi \rangle$ . [3]

### Question 3

Find the singularities for the following complex functions. In case there is a branch cut, show it by a suitable sketch. [2+2]

(a)  $\sec(z)$

(b)  $\log(z^2 - 3z + 2)$

### Question 4

Find the Greens function for an oscillator obeying:

$$\ddot{Q}(t) + \eta\dot{Q}(t) + \Omega^2 Q(t) = 0.$$

In the above,  $\dot{Q}(t) = \frac{dQ(t)}{dt}$ . [6]





## Question 5

(a) Consider two energy levels with energies  $\epsilon$  and  $-\epsilon$ . Total  $N$  classical distinguishable particles can occupy these levels, where the positions of these particles are fixed. If the total energy is  $E$ , write an expression for the number of configuration possible in terms of  $N$ ,  $E$  and  $\epsilon$ . [3]

(b) Considering  $N$ ,  $N_+$ ,  $N_- \gg 1$  (where  $N_{\pm}$  are number of particles in  $\pm\epsilon$  state), write an expression of the entropy in terms of  $N$ ,  $E$  and  $\epsilon$ . (Hint: for a large number  $p$ ,  $\ln(p!) \approx \ln(p) - p$ .) [4]

(c) Show that, the entropy from (b) is extensive ( $E$  is also extensive). [3]


## Question 6

A bead of mass  $m$  is constrained to move along a massless and frictionless rod ( $AB$ ) fixed at a constant angle  $\theta$  with the vertical axis as shown in the figure. The rod rotates with a uniform angular speed  $\omega$ . At a time  $t$ , the distance of the instantaneous position  $P$  of the bead from  $A$  is denoted by  $r$ . The distance  $OA$  is  $h$ .

(i) Set up the Lagrangian for the bead in terms of given variables assuming at  $t = 0$ , the rod is in  $XZ$  plane. (Direct writing of the expression will be awarded zero credit). [4]

(ii) Using Euler-Lagrange equation, find the equation of motion of the bead. [2]

(iii) Solve for  $r$  at any arbitrary time  $t$  assuming that the bead was initially at rest and was situated at  $A$ . [4]





## Question 7

(a) Write down the Maxwell equations for electrodynamics (in terms of  $\mathbf{E}$  and  $\mathbf{B}$  fields) in presence of free charges and free current. [ 1]

(b) Convert the inhomogeneous Maxwell equations in terms of the magnetic vector potential  $\mathbf{A}$  and the electromagnetic scalar potential  $\Phi$ . [ 4]

(c) From your last result write down the inhomogeneous Maxwell equations in the Lorenz gauge and the Coulomb gauge. Clearly state the Lorenz gauge condition and the Coulomb gauge condition. [ 2]

(d) Given a scalar and vector potential satisfying Lorenz gauge condition, find the transformations of these scalar and vector potential such that Lorenz gauge condition is still satisfied. [3]

## Question 8

List an experiment each that can be used for measuring the following fundamental constants in laboratory describing in brief (in two lines) the principle used in them. [ $2 \times 5 = 10$ ]

- (a) Electron charge ( $e$ )
- (b) Boltzmann constant ( $k_B$ )
- (c) Plank's constant ( $h$ )
- (d) speed of light ( $c$ )
- (e) permittivity of free space ( $\epsilon_0$ )

## Question 9


(a) Using IC-741, design a circuit which satisfies the following equation: [5]

$$V_{out} = \frac{dV_{in}}{dt} + 2V_{in} + 2$$





(b) For the following circuit, calculate the dc 'Q' point. [5]



Zero Vigyan